

On the Mean Parton Transverse Momentum versus Associated Hadron p_t in Di-Hadron Correlations at RHIC and LHC

A. Morsch

February 2, 2008

Abstract

From Pythia simulations of pp collisions we determine the mean transverse momentum $\langle p_t^{\text{parton}} \rangle$ of partons producing the leading and associated particles studied in di-hadron correlations. The dependence of $\langle p_t^{\text{parton}} \rangle$ on the associated particle transverse momentum and expected differences at RHIC and LHC centre of mass energies are discussed.

1 Introduction

Azimuthal correlations of high- p_t hadrons are used at RHIC to study the fragmentation properties of partons produced in $2 \rightarrow 2$ hard scatterings. The correlations are studied as a function of the transverse momentum of the trigger hadron p_t^{trig} and the associated particle p_t^{assoc} [1, 2]. The angular correlation functions show two characteristic approximately Gaussian peaks for hadrons produced from the fragmentation of the same parton that produced the trigger hadron (near-side) peak and from the fragmentation of the partons produced back-to-back approximately balancing the transverse momentum of the trigger parton (away-side peak). A strong suppression of the back-side peak in central Au+Au collisions relative to pp collisions has been observed and interpreted as evidence for in-medium partonic energy loss.

It has been demonstrated that due to the bias induced by the steeply falling production spectrum $d\sigma/dp_t^{\text{parton}}$, high- p_t hadrons carry about 60 – 70% of the original parton transverse momentum (see for example [3]). Due to the steepness of the partonic fragmentation function $D(z)$ with $z = p_t^{\text{assoc}}/p_t^{\text{parton}}$, it can be expected that demanding the presence of an additional associated particle further increases the parton mean transverse momentum $\langle p_t^{\text{parton}} \rangle$ ([4]): for constant p_t^{assoc} , z decreases with increasing p_t^{parton} . This effect should be stronger for the near-side associated hadrons, owing to the trivial fact that p_t^{parton} can not be smaller than the sum $p_t^{\text{trig}} + p_t^{\text{assoc}}$. The on average higher p_t^{parton} and the corresponding smaller production cross-section have to be taken

in account when comparing near-side to away-side associated hadron p_t spectra and widths of angular correlations.

In the next sections we show the results of Pythia simulations establishing the relation between p_t^{parton} and the associated parton transverse momentum in the range $p_t^{\text{assoc}} > 0 - 8 \text{ GeV}$ for $p_t^{\text{trig}} > 8 \text{ GeV}$ at RHIC and LHC energies. The choice of cuts may allow to use these results for further interpretation of the di-hadron correlation data recently published by the STAR collaboration [5].




Figure 1: Trigger normalised charged hadron fragmentation function $D(z_T)$ with $p_t^{\text{trig}} > 8 \text{ GeV}$ obtained from a Pythia simulation of pp collisions at $\sqrt{s} = 200 \text{ GeV}$ as described in the text. The simulation results are compared with the STAR d-Au data [5] (data values read from the figure).

2 Method

Pythia 6.2 [6, 8] is used to simulate pp collisions at $\sqrt{s} = 200 \text{ GeV}$ and 5500 GeV (QCD processes, MSEL=1). The transverse momenta of all final state charged hadrons and the partons resulting from the hard $2 \rightarrow 2$ scattering process are recorded. We select the hadron with the highest p_t , p_t^{max} within the pseudo rapidity window $|\eta| < 1$, the trigger hadron. If $p_t^{\text{max}} > p_t^{\text{trig}}$ the event is selected. Subsequently we identify the highest p_t^{assoc} hadron within a distance $\Delta\phi < 0.63$ to the trigger hadron, the associated near-side hadron and within $|\Delta\phi - \pi| < 0.63$, the associated away-side hadron.

The associated parton transverse momentum is determined by identifying the parton emerging from the hard scattering closest in $\eta - \phi$ to the trigger particle. Alternatively we used the Pythia PYCELL cone jet reconstruction algorithm (with $R = 1$) to find the jet with the axis closest to the trigger hadron. Both methods give similar results. The average parton momentum p_t^{parton} is determined for different $(p_t^{\text{trig}}, p_t^{\text{assoc}})$ -cuts.

As a benchmark for our simulations we show in Fig. 1 the $D(z_T)$ ($z_T = p_t^{\text{assoc}}/p_t^{\text{trig}}$) distributions for near-side and away-side correlated particles. The distributions agree very well with the STAR d-Au data [5].




Figure 2: The parton transverse momentum spectrum selected by $p_t^{\text{trig}} > 8 \text{ GeV}$, $p_t^{\text{assoc}} > 0 \text{ GeV}$ (solid line) is compared to the one selected by $p_t^{\text{trig}} > 8 \text{ GeV}$, $p_t^{\text{assoc}} > 2 \text{ GeV}$. (dashed line, online:red). The distribution corresponds to a statistics of 10^4 triggers.

3 Discussion of the results

Fig. 2 compares the parton spectra selected by $p_t^{\text{trig}} > 8 \text{ GeV}$ and $p_t^{\text{assoc}} > 0 \text{ GeV}$ to the one selected by $p_t^{\text{trig}} > 8 \text{ GeV}$ and $p_t^{\text{assoc}} > 2 \text{ GeV}$. Already at this moderate associated hadron p_t an important shift to higher p_t^{parton} and a sizable reduction of rate is observed.

Fig. 3 shows $\langle p_t^{\text{parton}} \rangle$ as a function of p_t^{assoc} . $\langle p_t^{\text{parton}} \rangle$ rises approximately linearly in the range $0 < p_t^{\text{assoc}} < p_t^{\text{trig}}$. At RHIC the difference in $\langle p_t^{\text{parton}} \rangle$ for near-side and away-side associated partons reaches up 30% at high p_t^{assoc} . At LHC energies the difference is about a factor of 1.5 smaller and $\langle p_t^{\text{parton}} \rangle$ is a factor of 1.5 higher. The difference is due to the different shape of the partonic production cross-section as a function of p_t^{parton} ; $d\sigma/dp_t^{\text{parton}}$ follows a power law $1/(p_T^{\text{parton}})^n$, where for parton energies considered here, $n \approx 8$ at RHIC and ≈ 4 at LHC.

In a recent paper [5] the STAR collaboration reports on the direct observation of dijets in central Au+Au collisions at $\sqrt{s_{NN}} = 200 \text{ GeV}$. The transverse momentum distributions of near- and away-side associated hadrons is used to study the effect of the medium on dijet fragmentation. The authors emphasize that $D(z_T)$ is measurable without direct knowledge of the parton energy. Although this is true, one has to keep in mind when comparing near-side and




Figure 3: Average parton transverse momentum (p_t^{parton}) as a function of the associated hadron transverse momentum for $p_t^{\text{trig}} > 8$ GeV in pp collisions at $\sqrt{s} = 200$ GeV and $\sqrt{s} = 5500$ GeV.

away-side associated hadrons, that z_T covers different parton energy regions.

One directly observable effect is the kinematic suppression of the nearside peak seen in the STAR d-Au data. For increasing p_t^{assoc} , the amplitude of the near-side peak decreases relative to the away-side peak. The azimuthal correlation functions for $p_t^{\text{trig}} > 8$ GeV and $p_t^{\text{assoc}} > 6$ GeV as obtained from the Pythia simulations are shown in Fig. 4. Comparing the amplitude of the peaks at $\sqrt{s} = 200$ GeV to $\sqrt{s} = 5500$ GeV one observes that the near-side peak rises by a factor of ≈ 7 , whereas the away-side peak decreases by a factor of ≈ 1.3 .

The rise of the near-side peak is mainly due to the difference in the partonic production spectra for the two different energies. Since the difference between near-side and away-side $\langle p_t^{\text{parton}} \rangle$ is smaller at LHC the rise of the away-side peak due to the same effect is smaller. In addition, there are at least two effects that are expected to lead to the suppression of the away-side peak at LHC energies relative to RHIC.

- Broadening of the pseudorapidity correlations due to the smaller Bjorken $x_B \sim 1/\sqrt{s}$ leads to a reduction by a factor of 2.4 for $|\eta| < 1$.
- Broadening of the azimuthal correlations due to the increased contribution from higher order QCD processes. The width of the away-side peak is by a factor of two larger at LHC energies.




Figure 4: Azimuthal correlation function for $p_t^{\text{trig}} > 8 \text{ GeV}$, $p_t^{\text{assoc}} > 6 \text{ GeV}$ and $|\eta| < 1$ for pp collisions at $\sqrt{s} = 200 \text{ GeV}$ (solid line) and $\sqrt{s} = 5500 \text{ GeV}$ (dashed line).

The increased nearside correlation at higher centre of mass energies can already be seen in data on charged particle correlations published by the CDF collaboration [10] for $\sqrt{s} = 1800 \text{ GeV}$ ¹.

The systematics of the ratios between near-side to away-side peaks for different p_t^{assoc} is obtained by comparing the corresponding fragmentation functions $D(z_T)$ (Fig. 5).

4 Conclusions

From Pythia simulations of pp collisions at RHIC and LHC energies we have determined the mean transverse momentum $\langle p_t^{\text{parton}} \rangle$ of partons fragmenting into a leading trigger hadron with transverse momentum p_t^{trig} and/or an associated hadron with transverse momentum p_t^{assoc} for $p_t^{\text{trig}} > 8 \text{ GeV}$ and $0 < p_t^{\text{assoc}} < p_t^{\text{trig}}$. For both, near-side and away-side correlations, p_t^{parton} depends not only on the p_t^{trig} but also strongly on the associated hadron p_t^{assoc} . Due to the additional kinematic constraint $p_t^{\text{parton}} > p_t^{\text{trig}} + p_t^{\text{assoc}}$, $\langle p_t^{\text{parton}} \rangle$ is higher

¹In simulation, the forward-backward peak inversion between RHIC and LHC energies has first been remarked in a HIJING study [9]




Figure 5: Trigger normalised charged hadron fragmentation function $D(z_T)$ with $p_t^{\text{trig}} > 8 \text{ GeV}$ obtained from a Pythia simulation of pp collisions at $\sqrt{s} = 200 \text{ GeV}$ (left) and $\sqrt{s} = 5500 \text{ GeV}$ (right).

for near-side correlated hadrons than for away-side correlations. The difference amounts to up to 30% at RHIC and 20% at LHC.

The higher $\langle p_t^{\text{parton}} \rangle$ and the corresponding smaller production cross-section have to be taken in account when comparing near-side to away-side associated hadron transverse momentum spectra and widths of angular correlations. In particular, at LHC energies, where the partonic production cross-section $d\sigma/dp_t^{\text{parton}}$ is less steep than at RHIC energies, the near-side azimuthal correlation peak will be enhanced with respect to the one observed at RHIC.

Acknowledgements

The author greatly appreciates the fruitful discussions with G. Paic.

References

- [1] C. Adler *et al.* (STAR Collaboration), Phys. Rev. Lett. **90** (2003) 082302.
- [2] S. S. Adler *et al.* (PHENIX Collaboration), Phys. Rev. Lett. **91** (2003) 072301.
- [3] X. N. Wang, Phys. Lett. B **579**, (2004) 299.
- [4] S. S. Adler *et al.* (PHENIX Collaboration), hep-ex/0605039.
- [5] J. Adams *et al.* (STAR collaboration), nucl-ex/0604018.
- [6] T. Sjostrand, Phys. Lett. B **157** (1985) 321.
- [7] M. Bengtsson, T. Sjostrand, and M. van Zijl, Z. Phys. C **32** (1986) 67.

- [8] T. Sjostrand and M. van Zijl, Phys. Rev. D**36**, (1987) 2019.
- [9] H. Appelshäuser and M. Ploskon, ALICE Internal Note ALICE-INT-2005-49.
- [10] T. Affolder *et al.* (CDF Collaboration) Phys. Rev. D**65** (2002) 092002.